

Beyond experimentation: Five trajectories of cigarette smoking in a longitudinal sample of youth

Lauren M. Dutra, Stanton A. Glantz, Nadra E. Lisha, Anna V. Song

Published: February 9, 2017 • <https://doi.org/10.1371/journal.pone.0171808>

Abstract

The first goal of this study was to identify the most appropriate measure of cigarette smoking for identifying unique smoking trajectories among adolescents; the second goal was to describe the resulting trajectories and their characteristics. Using 15 annual waves of smoking data in the National Longitudinal Survey of Youth 1997 (NLSY97), we conducted an exploratory latent class growth analysis to determine the best of four outcome variables for yearly smoking (cigarettes per day on days smoked, days smoked per month, mean cigarettes per day, and total cigarettes per month) among individuals aged 12 to 30 ($n = 8,791$). Days smoked per month was the best outcome variable for identifying unique longitudinal trajectories of smoking and characteristics of these trajectories that could be used to target different types of smokers for prevention and cessation. Objective statistics were used to identify four trajectories in addition to never smokers (34.1%): experimenters (13.6%), quitters (8.1%), early established smokers (39.0%), and late escalators (5.2%). We identified a quitter and late escalator class not identified in the only other comparable latent class growth analysis. Logistic regressions were used to identify the characteristics of individuals in each trajectory. Compared with never smokers, all trajectories except late escalators were less likely to be black; experimenters were more likely to be out of school and unemployed and drink alcohol in adolescence; quitters were more likely to have a mother with a high school degree/GED or higher (versus none) and to use substances in adolescence and less likely to have ever married as a young adult; early established smokers were more likely to have a mother with a high school diploma or GED, be out of school and unemployed, not live with both parents, have used substances, be depressed, and have peers who smoked in adolescence and to have children as young adults and less likely to be Hispanic and to have ever married as young adults; and late escalators were more likely to be Hispanic, drink alcohol, and break rules in adolescence and less likely to have ever married as young adults. Because of the number of waves of data analyzed, this analysis provided a clearer temporal depiction of smoking behavior and more easily distinguishable smoking trajectories than previous analyses. Tobacco control interventions need to move beyond youth-focused approaches to reach all smokers.

Citation: Dutra LM, Glantz SA, Lisha NE, Song AV (2017) Beyond experimentation: Five trajectories of cigarette smoking in a longitudinal sample of youth. PLoS ONE 12(2): e0171808. <https://doi.org/10.1371/journal.pone.0171808>

Editor: Olga Y. Gorlova, Dartmouth College Geisel School of Medicine, UNITED STATES

Received: May 25, 2016; **Accepted:** January 26, 2017; **Published:** February 9, 2017

Copyright: © 2017 Dutra et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: All relevant data are within the paper and its Supporting Information files.

Funding: This research was supported by R01 CA-060121 (Drs. Glantz and Song), R25CA-113710 and P50CA180890 (Dr. Dutra), 1U01CA154240 (Dr. Lisha) from the National Institutes of Health and Food and Drug Administration Center for Tobacco Products. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the US FDA. This research was conducted with Bureau of Labor Statistics data. The views expressed here do not necessarily reflect the views of the Bureau of Labor Statistics.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Tobacco-induced disease remains the leading preventable cause of death in the United States [1], with minorities bearing a disproportionate burden of the disease burden [2]. Because 80% of adult smokers begin smoking before age 18 [3], tobacco control efforts often focus on preventing adolescents from smoking their first cigarette. However, only one-third of youth who experiment with cigarettes ever become regular smokers [3,4]. As a result, these programs miss a key opportunity to prevent the transition from experimentation to established smoking, which may occur in the mid-to-late 20s [5]. In combination with growing recognition that young adulthood (ages 18 to 25) is a critical period of vulnerability [6], particularly due to significant life changes [3] such as starting college, separating smokers into different trajectories (patterns of smoking) and identifying when escalation and de-escalation occurs can inform efforts to prevent transition to regular smoking.

In addition to several regional U.S. latent class growth analyses of cigarette smoking dating back to the 1980s [4,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29], three national studies of the four-wave longitudinal AddHealth (years 0, 1, 5–6, and 13–14) demonstrated that multiple trajectories of smoking behavior exist [30,31,32]. Two of the national analyses followed participants for the first three waves (to age 25) and identified five trajectories in addition to never smokers [30,32]. The third, by Fuemmeler et al. [31], used all four waves to track smoking behavior into young adulthood (age 32) and identified four trajectories, in addition to never smokers. These national analyses identified characteristics that distinguished between trajectories, including alcohol or drug use, deviance, maternal smoking, peer smoking, conduct problems, depressive symptoms, and state prevalence of adolescent smoking. (Covariates included gender, race/ethnicity, and parental education.) These studies did not assess school enrollment status, which is essential to determining which types of smokers are exposed to school-based prevention and cessation programs. These studies were limited to four waves, with multiple years between assessments. In addition, these studies only examined baseline predictors of membership in a given trajectory of smoking behavior. As a result, they did not identify characteristics of high-risk smokers that could be used to target them in young adulthood. The fact that they used two different measures of smoking (mean cigarettes per day and total cigarettes per month) does not allow a decision to be made about the best measure of smoking for assessing youth and young adult smoking behavior, which is typically lighter and more intermittent than among established adult smokers [33,34]. This paper reports a model building process using latent class growth analysis based on 15 years of annual data from the National Longitudinal Survey of Youth 1997 (NLSY97) to determine the best measure of smoking to identify distinct behavioral trajectories of smoking and differentiate between youth and young adults who progress to established smoking and those who quit smoking. By identifying these trajectories and the adolescent and young adult characteristics of those who follow them, this is the first analysis that can be used to create targeted national public health and clinical interventions for smoking prevention and early cessation that extend through young adulthood.

Materials and methods

Sample

The U.S. Bureau of Labor Statistics NLSY97 [35] began in 1997 with a sample of 8,984 youth aged 12 to 16 and provides 15 years of annual data on smoking (ages 26 to 30 at wave 15). The NLSY includes a nationally representative sample ($n = 6,748$) and an oversample of Hispanic/Latino and black respondents ($n = 2,236$) divided into five cohorts based on age (12 to 16) on December 31, 1996. Eighty-three percent (7,423 of 8,984) of those sampled in 1997 remained in 2011.

The analysis, which is based on the unrestricted public use NLSY97 dataset, included the 8,791 (98%) NLSY97 respondents with at least three non-missing years of data for 30-day cigarette use because a minimum of three points is necessary for latent class growth analysis (LCGA) [36]. All analyses used Bureau of Labor Statistics baseline weights [37] to adjust for NLSY sampling techniques, including the oversample of racial/ethnic minorities. This is an analysis of a de-identified dataset and was ruled exempt by the University of California San Francisco Committee on Human Research.

Measures

Dependent variable: Smoking.

Because there is no agreed upon measure of smoking for latent class growth analyses of youth and young adult smoking [30,31,32] and in order to determine the best distribution of the outcome data [38,39], we evaluated four smoking variables used in previous analyses: mean cigarettes smoked per day, cigarettes per day on the days smoked, days smoked per month, and total cigarettes smoked per month [4,8,15,30,31,40,41].

Independent variables.

Race/ethnicity, gender, socioeconomic status (SES), and employment/school enrollment status at age 16 were examined as predictors of trajectory membership. Age 16 was chosen for childhood variables (such as employment/school enrollment status) because data were available for all respondents at this age and enrollment in school was mandatory. Race/ethnicity was coded as non-Hispanic white, non-Hispanic black, Hispanic, and non-Hispanic mixed race or non-Hispanic other. Gender was coded as male/female. Household poverty status at age 16 and highest level of biological mother's education were used to measure childhood SES. Household poverty status was coded in relation to the previous year's federal poverty line into four roughly equal categories (analyzed as continuous): below poverty line (0), poverty line to 199%, greater than poverty line (1), 200%–299% of the poverty line (2), and 300% or more of the poverty line (3). Biological mother's highest education (assessed in 1997) was coded categorically as no degree (0), GED/high school degree (1), undergraduate degree (associate/junior college or bachelor) (2), and graduate school or above (masters, PhD, or professional degrees, such as MD or JD) (3). Employment/school enrollment status at age 16, included to reflect engagement, was categorically coded as enrolled in school but unemployed (0), employed but not enrolled in school (1), enrolled in school and employed (2), or neither employed nor enrolled in school (3). Age in 1997 was included in growth models to adjust for cohort effects [42].

We also tested several individual and interpersonal variables that could not be coded by age but were important to examine because they were included in previous LCGAs [30,31,32] of the AddHealth dataset. Depression in 2000 was coded continuously by taking the mean of responses to the five-question Mental Health Inventory adapted for use in the NLSY97 [43], with higher values reflecting higher levels of depression. The questions, which all had a four-point response scale ranging from "all of the time" (coded as 0) to "none of the time," (coded as 3) included "How much of the time during the last month have you..." "been a nervous person," "felt calm and peaceful," "felt down or blue," "been a happy person," and "felt so down in the dumps that nothing could cheer you up?" Peer smoking in 1997, assessed by the question "What percentage of kids [in your grade/in your grade when you were last in school] [smoke/smoked] cigarettes?," was coded continuously so that higher values of the variable reflected higher peer smoking using the five-point response scale that ranged from "almost none (less than 10%)" (coded as 0) to "almost all (more than 90%)" (coded as 4). We also measured conduct problems/rebellion in 2008, assessed using the question, "When I was in school, I used to break rules quite regularly," as a continuous variable based on the 7-point response scale ranging from "disagreed strongly" (0) to "agree strongly" (6), with higher values reflecting a greater tendency for rule-breaking. We also assessed substance use in 1997 via dichotomized ever use of alcohol, marijuana, and "cocaine or other hard drugs" (reference category was never

use). We assessed childhood family stability by dichotomizing responses to a question about whether the respondent lived with both biological parents in 1997, with a value of “1” signifying an individual who did not live with both biological parents in 1997 (and 0 as reference).

We examined several sociodemographic characteristics at age 26, the oldest available age for all respondents, to identify young adult characteristics of the different smoking trajectories. Marital status was ever married (married, separated, divorced, or widowed) versus never married (reference). Number of children was coded as one or more versus none (reference). Respondent’s highest level of education at age 26 was coded similarly to mother’s education.

Analysis

LCGA is a person-centered approach to modeling that can be used to identify individuals with similar behavioral trajectories in adolescence and young adulthood [44]. Never smokers (34.1% of sample, $n = 3,147$), those who reported 0 days smoked in the past 30 days for all non-missing years of data collection, were specified as an *a priori* class and excluded from the LCGA [14]. This established approach [45] reduced the computational burden of the large number of zeros (from never smokers) and provided a better fit to the distribution.

We used *Mplus* Version 7.3 because it accounts for the longitudinal structure of the NLSY and for missing data using full information maximum likelihood [46].

Determining the outcome variable.

We first conducted a model building process to determine the best smoking variable to achieve the goal of LCGM, to produce “meaningful patterns that distinguish subgroups of people” [30] that could be targeted by public health interventions. We first constructed single-class LCGMs in order to make sure all models converged [47]. For mean cigarettes per day, we used a linear LCGM because the outcome was continuous. For all other measure of smoking, we used negative binomial LCGMs because they accommodate count variables and, unlike the Poisson distribution, account for unobserved heterogeneity in observations, leading to more accurate estimation of standard errors [48].

Because traditional fit statistics [49] were unavailable in *Mplus* because we used count variables [50] and had more than eight time points [51], to provide further guidance we examined the shape of the trajectories starting with a 4-class LCGM [31]. This model building process is an established method [4,8,15,31,47] to identify covariates, shapes of trajectories, and relationships between outcome and explanatory variables. [38,40,41]

Determining the number of trajectories.

After we established the measure of smoking that provided the clearest differentiation between trajectories, a combination of theory and fit statistics was used to determine the ideal number of trajectories, starting with four [52]. The Bayesian Information Criteria (BIC), the Vuong-Lo-Mendell-Rubin Likelihood Ratio Test (LMR-LRT), the posterior probabilities, and the fraction of smokers in the trajectories were used to determine the most parsimonious model [47]. The BIC is a log-likelihood based statistic that accounts for number of model parameters, with lower values indicating better fit [52]. The LMR-LRT is a goodness-of-fit test that quantifies the likelihood that data can be described by a model with one less trajectory; a significant p-value (<0.05) indicates that the model with one additional trajectory has superior fit [52]. Respondents were assigned to the trajectory with the highest posterior probability, with probabilities above 0.7 considered acceptable [53]; all respondents met this criteria for trajectory assignment.

A cohort-sequential design [30] was used to link cohorts by age, resulting in 19 years (15 waves) of data (ages 12 to 30). This approach accounts for age-based cohort effects by designating age, not calendar year, as the time variable. We confirmed the validity of this approach by running separate LCGAs for each age group (12, 13, 14, 15, and 16 years old at baseline), which produced similar patterns of smoking at similar ages.

Determining predictors of trajectory membership.

After the outcome variable and number of trajectories were determined and respondents were assigned to trajectories by *Mplus*, Holm-Sidak adjusted chi-square and t-tests and logistic regressions (never smokers were the reference group to make our results comparable to those of previous LCGAs of the AddHealth national samples [30,31,32]) were used to examine differences in participant characteristics across trajectories [54]. All analyses (bivariate tests and logistic regressions) were completed using Stata version 12 (StataCorp) with the *svyset* command to adjust for weights. Respondent’s highest education was not included in adjusted models because of collinearity with employment/school enrollment status and mother’s education. We included youth SES measures (income and mother’s education) and educational engagement (employment/school enrollment status) because these factors are more stable over time than adult income and education. The final model, which included age in 1997 to adjust for age-cohort effects, included all predictor variables that were significant in bivariate analyses. We also examined prevalence of daily smoking by trajectory, median years of daily smoking, and median years between first reported smoking and daily smoking (adjusted for weights) to further understand patterns of smoking in each trajectory.

Results

All of the univariate latent class growth models converged except for mean cigarettes per day; therefore, this outcome was discarded. We then compared the trajectories of the other three outcome variables (S1 Table). We chose days per month as our outcome variable because it produced the most unique trajectories and a visible differentiation between smokers who progressed to daily smoking versus those who quit [30]. Days smoked per month is also a preferable measure of smoking intensity for youth and young adults because this variable provided unique patterns of smoking behavior and because cigarettes per day can be unstable over time [55]. Choosing this variable also achieved our aim of capturing progression (or lack thereof) to regular smoking and intermittent and nondaily smoking, which is a common trend among adolescents and young adults [33,34].

Overall sample characteristics

Approximately half of the sample was female (48.7%) and was primarily non-Hispanic white (66.6%) (Table 1). About half the respondents (52.9%) had mothers whose highest education was GED/high school diploma. At age 16, 88.6% were in school, including 54.7% who were employed while attending school. Mean household income was about twice (mean = 192%, median = 200%) the poverty level. In 1997, mean peer smoking was 1.63 (on average, participants estimated that approximately between 25% and 50% of their peers smoked), mean likelihood of breaking rules in school was 2.26 (between “disagree a little” and “neither agree nor disagree”), 47.3% did not live with both biological parents in 1997, 45.1% had ever drunk alcohol, 21.0% had ever tried marijuana, and 7.1% had ever used cocaine or hard drugs in 1997. Mean depression score in 2000 (range from 0 to 4) was 0.93 (symptoms of depression “some of the time”).

Characteristics	Overall sample	Experimenters		Quitters		Early established smokers		Late escalators	
	(n = 7,972)	(n = 1,205)	(n = 701)	(n = 3,205)	(n = 533)	(n = 3,205)	(n = 533)	(n = 3,205)	(n = 533)
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
Demographics									
Female	48.7%	48.7%	48.7%	48.7%	48.7%	48.7%	48.7%	48.7%	48.7%
Hispanic	13.0%	13.0%	13.0%	13.0%	13.0%	13.0%	13.0%	13.0%	13.0%
Non-Hispanic white	66.6%	66.6%	66.6%	66.6%	66.6%	66.6%	66.6%	66.6%	66.6%
Non-Hispanic black	25.2%	25.2%	25.2%	25.2%	25.2%	25.2%	25.2%	25.2%	25.2%
Non-Hispanic other	8.2%	8.2%	8.2%	8.2%	8.2%	8.2%	8.2%	8.2%	8.2%
Mother's highest education									
1 - Less than high school	12.8%	12.8%	12.8%	12.8%	12.8%	12.8%	12.8%	12.8%	12.8%
2 - GED/high school diploma	52.9%	52.9%	52.9%	52.9%	52.9%	52.9%	52.9%	52.9%	52.9%
3 - Associate's degree	27.8%	27.8%	27.8%	27.8%	27.8%	27.8%	27.8%	27.8%	27.8%
4 - Bachelor's degree	7.2%	7.2%	7.2%	7.2%	7.2%	7.2%	7.2%	7.2%	7.2%
5 - Graduate professional degree	0.5%	0.5%	0.5%	0.5%	0.5%	0.5%	0.5%	0.5%	0.5%
Family and household characteristics									
Family size (age 16)									
1 - 1 child, non-employed	33.8%	33.8%	33.8%	33.8%	33.8%	33.8%	33.8%	33.8%	33.8%
2 - 1 employed, non-employed	4.1%	4.1%	4.1%	4.1%	4.1%	4.1%	4.1%	4.1%	4.1%
3 - 2 employed, non-employed	1.7%	1.7%	1.7%	1.7%	1.7%	1.7%	1.7%	1.7%	1.7%
4 - 3 employed, non-employed	0.6%	0.6%	0.6%	0.6%	0.6%	0.6%	0.6%	0.6%	0.6%
5 - 4 employed, non-employed	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%
6 - 5 employed, non-employed	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%
Household income (age 16)	1.92 (1.41)	2.00 (1.41)	1.92 (1.41)	2.00 (1.41)	1.92 (1.41)	2.00 (1.41)	1.92 (1.41)	2.00 (1.41)	1.92 (1.41)
Household income (2000)	0.93 (0.80)	0.93 (0.80)	0.93 (0.80)	0.93 (0.80)	0.93 (0.80)	0.93 (0.80)	0.93 (0.80)	0.93 (0.80)	0.93 (0.80)
Peer smoking	1.63 (1.00)	1.63 (1.00)	1.63 (1.00)	1.63 (1.00)	1.63 (1.00)	1.63 (1.00)	1.63 (1.00)	1.63 (1.00)	1.63 (1.00)
Breaks rules in school (2000)	2.26 (1.20)	2.26 (1.20)	2.26 (1.20)	2.26 (1.20)	2.26 (1.20)	2.26 (1.20)	2.26 (1.20)	2.26 (1.20)	2.26 (1.20)
Ever married	55.3%	55.3%	55.3%	55.3%	55.3%	55.3%	55.3%	55.3%	55.3%
Ever had children	54.4%	54.4%	54.4%	54.4%	54.4%	54.4%	54.4%	54.4%	54.4%
Household gender parity	47.4%	47.4%	47.4%	47.4%	47.4%	47.4%	47.4%	47.4%	47.4%
Ever used alcohol	45.1%	45.1%	45.1%	45.1%	45.1%	45.1%	45.1%	45.1%	45.1%
Ever used marijuana	21.0%	21.0%	21.0%	21.0%	21.0%	21.0%	21.0%	21.0%	21.0%
Ever used cocaine/hard drugs (1997)	7.1%	7.1%	7.1%	7.1%	7.1%	7.1%	7.1%	7.1%	7.1%
Adult variables (age 30)									
Ever smoked	49.5%	49.5%	49.5%	49.5%	49.5%	49.5%	49.5%	49.5%	49.5%
Median years of daily smoking	0 (0-3)	6 (1-12)	0 (0-2)	0 (0-2)	0 (0-2)	0 (0-2)	0 (0-2)	0 (0-2)	0 (0-2)
Median years of highest education	8.4%	8.4%	8.4%	8.4%	8.4%	8.4%	8.4%	8.4%	8.4%
1 - Less than high school	10.2%	10.2%	10.2%	10.2%	10.2%	10.2%	10.2%	10.2%	10.2%
2 - GED/high school diploma	55.2%	55.2%	55.2%	55.2%	55.2%	55.2%	55.2%	55.2%	55.2%
3 - Associate's degree	30.8%	30.8%	30.8%	30.8%	30.8%	30.8%	30.8%	30.8%	30.8%
4 - Bachelor's degree	4.7%	4.7%	4.7%	4.7%	4.7%	4.7%	4.7%	4.7%	4.7%
5 - Graduate professional degree	4.7%	4.7%	4.7%	4.7%	4.7%	4.7%	4.7%	4.7%	4.7%

Table 1. Individual-level characteristics by trajectory of cigarette smoking.
<https://doi.org/10.1371/journal.pone.0171808.t001>

By age 26, most of the respondents (59.3%) had never been married and had no children (54.4%). At age 26, 9.4% had no degree, 55.3% had a GED/high school diploma, 30.6% had a college degree, and 4.7% had a graduate or professional degree.

Trajectories of smoking from 1997–2011

We tested one to five trajectories. The BIC consistently declined across solutions (S2 Table). The LMR-LRT p-value increased across trajectories, reaching non-significance at five trajectories. Since this result suggested five trajectories was not superior to four and all posterior probabilities were over 0.7 for all solutions (0.90–0.98), we selected four classes.

Based on trajectory shapes and the literature, we named the four trajectories (in addition to never smokers, trajectory 0) experimenters (trajectory 1), quitters (trajectory 2), early established smokers (trajectory 3), and late escalators (trajectory 4) (Fig 1). Never smokers comprised 34.1% (n = 3,147) of the sample, experimenters 13.6% (n = 1,205), quitters 8.1% (n = 701), early established smokers 39.0% (n = 3,205), and late escalators 5.2% (n = 533).




Fig 1. Four trajectories of smoking behavior among ever smokers in the National Longitudinal Survey of Youth 1997 (NLSY97).
 Analyses of 19 years (15 waves) of smoking data (days smoked per month) for participants in the NLSY97 revealed four patterns of smoking behavior (in addition to never smokers): experimenters, quitters, early established smokers, and late escalators.
<https://doi.org/10.1371/journal.pone.0171808.g001>

Days smoked per month peaked at age 16 for experimenters (mean = 1.22; 95% CI: 0.96–1.48), with mean days smoked per month remaining below one for 16 of the 19 years. For quitters, days smoked peaked at age 17 (10.64; 9.61–11.67) and decreased to less than one day per month at age 25. For early established smokers, days smoked peaked at age 22 (22.60; 22.14–23.05) and plateaued by age 30 (18.49; 17.21–19.78). Late escalators' highest smoking was 9.00 (6.11–11.89) days per month at age 30 and still trending upward.

Only 2.0% of experimenters, 49.5% of quitters, 93.5% of early established smokers, and 24.0% of late escalators ever smoked daily. Median years of daily smoking was 0 for experimenters (10th–90th percentile: 0–0; only 2.0% ever smoked daily), 0 for quitters (0–3), 6 for early established smokers (1–12), and 0 for late escalators (0–2). Median years from first cigarette to daily smoking was 0 (0–1) for the 2.0% (23 of 1,205) of experimenters who made this transition, 1 year (0–4) for the 48.4% (339 of 701) of quitters who made this transition, 1 year (0–5) for the 92.5% (2,966 of 3,205) of early established smokers who made this

transition, and 3 years (0–11) for the 26.3% (140 of 533) of late escalators who made this transition. Median age of transition to daily smoking (for those who transitioned) was 15 (13–16) for experimenters, 17 (15–20) for quitters, 17 (14–21) for early established smokers, and 26 (24–28) for late escalators.

Predictors of trajectory membership

Our adjusted model included all covariates (Table 2). Compared with never smokers, experimenters were more likely to have ever tried alcohol in 1997 and less likely to be non-Hispanic black (than non-Hispanic white) and to be employed and enrolled in school at age 16 (versus not enrolled in school and unemployed). Quitters were more likely to be unemployed at age 16, to have a mother who had at least completed high school/GED, and to have ever drunk alcohol or used marijuana in 1997. They were less likely to be black, to be enrolled in school at age 16, and to have ever been married by age 26. Early established smokers were more likely to have a mother whose highest level of education was a GED/high school diploma (versus no degree), to have used alcohol or marijuana in 1997, have peers who smoked in 1997, to be depressed in 2000, have broken rules in school in 2008, and to have children at age 26. They were less likely to be black or Hispanic, to be enrolled in school or employed at age 16, to live with both biological parents in 1997, and to have ever been married at age 26. Late escalators were more likely to be Hispanic, to have drunk alcohol in 1997, and to have broken rules in school and less likely to have ever been married at age 26 than never smokers.

	Experimenters (trajectory 1) versus never smokers (trajectory 0)	Quitters (trajectory 2) versus never smokers (trajectory 0)	Early established smokers (trajectory 3) versus never smokers (trajectory 0)	Late escalators (trajecto ry 4) versus never smokers (trajectory 0)
Each variable's significance unless otherwise noted.				
Gender				
- Female	REF	REF	REF	REF
+ Male	1.20 (0.72–2.00)	1.61 (0.95–1.84)	1.40 (0.95–2.07)	1.98 (0.93–2.11)
Race/ethnicity				
- Non-Hispanic white	REF	REF	REF	REF
+ Non-Hispanic black	0.45 (0.24–0.82)	0.19 (0.09–0.40)	0.28 (0.18–0.45)	0.80 (0.37–1.75)
+ Hispanic	1.61 (0.67–3.89)	0.77 (0.33–1.81)	0.40 (0.24–0.69)	2.62 (1.14–6.07)
+ Non-Hispanic msaal	1.36 (0.43–4.31)	0.52 (0.05–1.92)	1.04 (0.38–2.82)	1.80 (0.99)
Mother's highest education				
- <GED/HS diploma	REF	REF	REF	REF
+ GED/HS graduate	1.68 (0.91–3.16)	3.29 (1.41–7.65)	1.92 (1.15–3.20)	2.94 (0.89–4.68)
+ A.A./BA/BS	1.62 (0.79–3.36)	2.92 (0.97–9.11)	1.50 (0.59–3.86)	2.70 (0.99–7.40)
+ Graduate/professional	2.07 (0.71–6.06)	4.24 (1.09–16.43)	0.83 (0.33–2.11)	2.10 (0.39–12.24)
Employment/school enrollment status (age 16)				
- In school, not employed	REF	REF	REF	REF
+ Employed, not in school	0.87 (0.43–1.76)	0.82 (0.35–2.40)	1.63 (0.85–3.13)	0.80 (0.29–2.30)
+ Employed and in school	1.41 (0.80–2.49)	0.79 (0.34–1.45)	0.68 (0.41–1.02)	0.88 (0.44–1.78)
+ Neither in school nor employed	2.52 (1.09–6.83)	2.80 (0.98–8.02)	4.34 (2.25–8.37)	0.77 (0.24–2.52)
Non-parent family	0.78 (0.49–1.28)	1.38 (0.77–2.47)	1.85 (1.12–3.42)	1.01 (0.51–1.97)
Ever drunk alcohol	2.00 (0.91–4.14)	2.08 (1.08–4.01)	2.00 (0.85–4.80)	2.21 (1.04–4.67)
Ever used marijuana	1.19 (0.63–2.25)	3.45 (1.70–6.99)	2.11 (0.97–5.03)	0.74 (0.28–1.97)
Ever used cocaine/hard drugs (1997)	0.80 (0.24–2.62)	1.01 (0.34–2.96)	1.57 (0.69–3.58)	1.00 (0.99)
Household income (age 16)	1.30 (0.69–1.71)	0.93 (0.60–1.30)	0.85 (0.60–1.06)	0.81 (0.55–1.17)
Age	0.78 (0.63–0.95)	0.79 (0.63–1.01)	0.85 (0.75–1.00)	0.87 (0.73–1.03)
Depression (2000)	1.17 (0.76–1.82)	1.45 (0.89–2.47)	1.70 (1.07–2.61)	1.21 (0.65–2.26)
Peer smoking	1.19 (0.86–1.66)	1.19 (0.80–1.83)	1.26 (1.08–1.48)	0.98 (0.75–1.27)
Living in non-parent rules in school (2008, retrospective)	1.08 (0.64–1.19)	1.15 (0.69–1.93)	1.28 (1.17–1.41)	1.28 (1.07–1.47)
Adult variables (age 26)				
Marital status				
- Never married	REF	REF	REF	REF
+ Ever Married	0.78 (0.49–1.29)	0.52 (0.28–0.94)	0.47 (0.31–0.70)	0.42 (0.25–0.81)
Has 1 child				
- No	REF	REF	REF	REF
+ Yes	1.28 (0.77–2.13)	1.12 (0.59–2.15)	1.76 (1.15–2.68)	1.10 (0.54–2.06)

Note: All models include all listed variables.
All models adjusted for stratification variables (to adjust for sampling techniques) and weights (to adjust for non-response and non-representativeness of sample) provided by the NLSY.
doi:10.1371/journal.pone.0171808.t002

Table 2. Adjusted odds of trajectory membership for all significant covariates (with 95% confidence intervals).
<https://doi.org/10.1371/journal.pone.0171808.t002>

All of the variance inflation factors were 1.44 or under in the final adjusted model, indicating little multicollinearity. The results of the unadjusted analyses (see Table 1) were similar to the adjusted results (see Table 2). The only variables that were significant in bivariate analyses but not significant in adjusted models were household income, which was significantly lower for early established smokers than members of the other trajectories (except late escalators), and ever use of cocaine/hard drug use, which was more common for experimenters and quitters than other trajectories.

Discussion

Exploratory analyses revealed that days smoked per month provided clearer distinction between trajectories of smoking than cigarettes per day on days smoked, mean cigarettes per day, or total cigarettes smoked per month. LCGA revealed four trajectories in addition to never smokers: experimenters, quitters, early established smokers, and late escalators. This finding contradicts the assumption that once adolescents initiate smoking, they all exhibit the same pattern of smoking: 21.4% of ever smokers were experimenters who never went on to become established smokers.

Trajectory characteristics

We found several qualitative and quantitative differences between our trajectories and those identified in LCGAs of AddHealth (S3 and S4 Tables) [30,31,32]. Fuemmeler et al. [31] did not identify a quitter class; this group is key to understanding why some individuals spontaneously quit. In addition, neither Fuemmeler et al. [31] nor Pollard et al. [32] identified a true late escalator class (started smoking after age 20 and trending upward at end of follow-up), thereby failing to identify an important minority group that has been ignored in the past. Fuemmeler et al. [31] and Pollard et al.'s [32] experimenter classes peaked much later (ages 24 and 25, respectively) than ours (age 16). Costello et al.'s [30] quitter class peaked at age 20, later than our quitter class, which peaked at age 16.

Some of these differences may be due to our specification of never smokers as an a priori class (as opposed to including never smokers in the LCGA). As a result, our never smoker category was smaller than those identified in analyses of AddHealth (S4 Table). In addition, our use of days smoked per month (versus cigarettes per day or a combined measure of frequency and intensity) distinguishes our analysis from those completed with AddHealth. Most importantly, the great number of waves of data collection in the NLSY97 (annually between 1997 and 2011) resulted in a level of detail unavailable when using AddHealth's three or four waves of data (1995, 1996, 2001–2002, 2008–2009). This important difference enabled us to identify specific ages at which key transitions occur.

For example, *early established* smokers escalated between age 12 and 22, *late escalators* escalated between age 22 and 30, and *quitters* de-escalated between 18 and 24. As a result, we concluded that late escalators should be targeted in young adulthood. In the Fuemmeler et al. [31] analysis, in particular, with the exception of the late heavy user group, the most distinguishing graphical characteristic between the classes was the intercept. In contrast, our analysis revealed trajectories with differing intercepts, slopes, and shapes.

Like the current analysis and analyses of Add Health [30,31,32], LCGAs of smoking of the sub-national samples [4,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29] provides support for the existence of multiple smoking trajectories. The findings of the sub-national studies, however, are less likely to be generalizable to the entire United States, so have limited applicability to the development of national policies and interventions for smoking prevention and cessation.

Characteristics of individuals in trajectories

Analyses of AddHealth [30,31,32] revealed that alcohol or drug use, deviance, maternal smoking, peer smoking, conduct problems, depressive symptoms, and state prevalence of adolescent smoking distinguished one or more smoking trajectories from never smokers. With the exception of state smoking prevalence, which we did not include in our analysis, our results demonstrated similar findings for these variables.

We also identified several characteristics that were not identified in previous analyses (S4 Table). Members of all trajectories except experimenters were less likely to have been married at age 26 compared with never smokers. Compared with never smokers, *experimenters* were less likely to be black and less likely to be enrolled in school or working at age 16. However, overall, experimenters were similar to never smokers. Our results suggest that tobacco control programs should expand their efforts beyond preventing experimentation to also preventing escalation to established smoking.

Quitters were less likely to be black, tended to have mothers with higher education, and were more likely to have ever used marijuana. Compared with early established smokers, quitters were more likely to be Hispanic and less likely to break rules in school. Additional research should identify further characteristics about this group and their reasons for quitting. However, the existence of this group likely indicates that traditional youth-based smoking prevention and cessation programs are successful for some adolescents.

Early established smokers were less likely to be black or Hispanic, to have educated mothers, and to grow up in a two-parent household; they were more likely to have ever used marijuana and to drop out of school by age 16. As a result, they are probably less likely to be reached by school-based interventions in adolescence.

Late escalators only exhibited some traditional smoking risk factors, were more likely to be Hispanic, and were still enrolled in school at age 16. Because their smoking escalates after age 18, they are more likely to benefit from interventions that extend into young adulthood.

Public health policy implications

This is the first analysis that can be used to create targeted national public health and clinical interventions for smoking prevention and early cessation that extend through young adulthood. Population-level interventions, including tobacco taxes, smokefree laws (particularly workplace) and rules (such as car), smokefree housing, and media campaigns are key to reaching all types of smokers [56,57,58]. People covered by smokefree laws are more likely to have smokefree cars [56] and homes [59], which increase the likelihood of successful smoking cessation [57,60]. In addition, as recommended by the American Academy of Pediatrics [61], all movies and video games depicting smoking should be given an R rating [62,63] or eliminated entirely. Similarly, restricting *all* tobacco advertising, including point-of-sale and product placement, would deter smoking initiation and adolescent smoking and progression to established smoking. In addition to these approaches, we identified specific methods of reaching early established smokers and late escalators. Because early established smokers are more likely to have children, one method of reaching them is by incorporating smoking cessation interventions into early intervention programs, such as Women, Infants, and Children, an assistance program for pregnant women and children under age 5, and California's "First Five," a county-based early intervention program [40], which is funded by tobacco taxes. To reach late escalators in young adulthood when their smoking frequency is increasing, we recommend smokefree schools, smokefree college campuses, college-based smoking cessation programs, and bar-based smoking cessation and prevention interventions [64].

The Truth Initiative, which targets youth and young adults [65], and the Centers for Disease Control and Prevention's *Tips From Former Smokers* media campaign, which targets 18- to 54-year-olds [66], are likely to be effective for early established smokers and late escalators. Media campaigns, such as the Food and Drug Administration's *The Real Cost* and *Fresh Empire* campaigns [41,67], should be extended to include individuals into their late 20s, particularly since youth-centric programs are an industry tactic [68].

Because of the immense gains made in tobacco control in the past 50 years, including denormalizing tobacco and revolutionizing social norms, it is necessary to reevaluate the efficacy of the established approach to tobacco prevention, particularly as new databases emerge. The Population Assessment of Tobacco and Health (PATH) study should be used to continue to identify patterns of smoking to prevent progression to established smoking and encourage smoking cessation and identify the effects of rising dual use of tobacco products (including e-cigarettes) and marijuana on trajectories of cigarette smoking.

Limitations

The primary limitation of this analysis is that it does not include policy variables that affect smoking. It is possible that individuals shift trajectories over time, but LCGAs do not accommodate such shifts.

Conclusion

Days smoked per month is a useful measure of smoking for identifying different types of smokers and should therefore be included in tobacco use surveys. Also, multiple patterns of smoking are necessary to fully describe adolescent and young adult smoking uptake and patterns of use. Just as the tobacco industry identified different types of smokers (market segments) to target, tobacco control programs can use information about types of smokers to develop more effective interventions [69]. Targeting young adults (18 to 24) is important because this group is most likely to quit and do so successfully. Pediatricians should collect information on smoking intensity and frequency and attempt to prevent smoking escalation among light smokers by helping them identify triggers, referring them to counseling resources, and encouraging youth to advocate a smokefree lifestyle for parents and peers. Because smoking is still dynamic beyond age 18, pediatricians and other physicians should ensure that information about tobacco use is transferred to general medicine practitioners as their patients get older. Our analysis also supports increasing the smoking age from 18 to 21 [61]; 16.9% (954) of the ever smokers in our analysis initiated smoking between the ages of 18 and 21.

Supporting information

S1 Table. Comparison of four-class solutions by smoking variable.

<https://doi.org/10.1371/journal.pone.0171808.s001>
(DOCX)

S2 Table. Fit statistics and sample size (N) by number of classes.

<https://doi.org/10.1371/journal.pone.0171808.s002>
(DOCX)

S3 Table. Graphical comparison of trajectories from other latent class growth analyses using national samples.

<https://doi.org/10.1371/journal.pone.0171808.s003>
(DOCX)

S4 Table. Comparison of this paper with comparable trajectories identified in previous latent class growth analyses of national datasets.

<https://doi.org/10.1371/journal.pone.0171808.s004>
(DOCX)

S1 Data. Raw data file as stata 12 DTA file.

<https://doi.org/10.1371/journal.pone.0171808.s005>
(DTA)

Acknowledgments

This research was conducted with restricted access to Bureau of Labor Statistics data. The views expressed here do not necessarily reflect the views of the Bureau of Labor Statistics.

Author Contributions

Conceptualization: SAG AVS.

Data curation: SAG.

Formal analysis: AVS NEL LMD.

Funding acquisition: SAG.

Investigation: LMD.

Methodology: SAG AVS NEL LMD.

Project administration: LMD SAG.

Resources: SAG.

Supervision: SAG AVS.

Validation: NEL LMD.

Visualization: SAG LMD.

Writing – original draft: LMD.

Writing – review & editing: SAG LMD AVS NEL.

References

1. Centers for Disease Control and Prevention (2012) Current Cigarette Smoking among Adults- United States, 2011. Morbidity and Mortality Weekly Report 61: 889–894. pmid:23134971
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

2. Centers for Disease Control and Prevention (2010) Racial/Ethnic Disparities and Geographic Differences in Lung Cancer Incidence—38 States and the District of Columbia, 1998–2006. *Morbidity and Mortality Weekly Report* 59: 1434–1438. pmid:21063273
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
3. U.S. Department of Health and Human Services (2012) Preventing Tobacco Use among Youth and Young Adults: A Report of the Surgeon General. Rockville, MD.
4. Chassin L, Presson CC, Pitts SC, Sherman SJ (2000) The Natural History of Cigarette Smoking from Adolescence to Adulthood in a Midwestern Community Sample: Multiple Trajectories and Their Psychosocial Correlates. *Health Psychology* 19: 223–231. pmid:10868766
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
5. Chen K, Kandel DB (1995) The Natural History of Drug Use from Adolescence to the Mid-Thirties in a General Population Sample. *American Journal of Public Health* 85: 41–47. pmid:7832260
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
6. Arnett JJ (2000) Emerging Adulthood: A Theory of Development from the Late Teens through the Twenties. *American Psychologist* 55: 469. pmid:10842426
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
7. Riggs NR, Chou CP, Li C, Pentz MA (2007) Adolescent to Emerging Adulthood Smoking Trajectories: When Do Smoking Trajectories Diverge, and Do They Predict Early Adulthood Nicotine Dependence? *Nicotine & Tobacco Research* 9: 1147–1154.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
8. White HR, Pandina RJ, Chen P-H (2002) Developmental Trajectories of Cigarette Use from Early Adolescence into Young Adulthood. *Drug and Alcohol Dependence* 65: 167–178. pmid:11772478
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
9. Brook JS, Zhang C, Burke L, Brook DW (2014) Trajectories of Cigarette Smoking from Adolescence to Adulthood as Predictors of Unemployment Status. *Nicotine & Tobacco Research* 16: 1559–1566.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
10. Mays D, Gilman SE, Rende R, Luta G, Tercyak KP, Niaura RS (2014) Parental Smoking Exposure and Adolescent Smoking Trajectories. *Pediatrics* 133: 983–991. pmid:24819567
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
11. Lee JY, Brook JS, Finch SJ, Brook DW (2016) Trajectories of Cigarette Smoking Beginning in Adolescence Predict Insomnia in the Mid Thirties. *Substance Use & Misuse* 51: 616–624.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
12. Brook JS, Pahl K, Ning Y (2006) Peer and Parental Influences on Longitudinal Trajectories of Smoking among African Americans and Puerto Ricans. *Nicotine & Tobacco Research* 8: 639–651.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
13. Karp I, O'Loughlin J, Paradis G, Hanley J, Difranza J (2005) Smoking Trajectories of Adolescent Novice Smokers in a Longitudinal Study of Tobacco Use. *Annals of Epidemiology* 15: 445–452. pmid:15967392
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
14. Orlando M, Tucker JS, Ellickson PL, Klein DJ (2004) Developmental Trajectories of Cigarette Smoking and Their Correlates from Early Adolescence to Young Adulthood. *Journal of Consulting and Clinical Psychology* 72: 400–410. pmid:15279524
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
15. Stanton WR, Flay BR, Colder CR, Mehta P (2004) Identifying and Predicting Adolescent Smokers' Developmental Trajectories. *Nicotine & Tobacco Research* 6: 843–852.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
16. Akhtar-Khaleel WZ, Cook RL, Shoptaw S, Surkan PJ, Teplin LA, Stall R, et al. (2016) Long-Term Cigarette Smoking Trajectories among Hiv-Seropositive and Seronegative Msm in the Multicenter Aids Cohort Study. *AIDS Behavior*.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
17. Orpinas P, Lacy B, Nahapetyan L, Dube SR, Song X (2016) Cigarette Smoking Trajectories from Sixth to Twelfth Grade: Associated Substance Use and High School Dropout. *Nicotine & Tobacco Research* 18: 156–162.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

18. Hampson SE, Tildesley E, Andrews JA, Barckley M, Peterson M (2013) Smoking Trajectories across High School: Sensation Seeking and Hookah Use. *Nicotine & Tobacco Research* 15: 1400–1408.
View Article • PubMed/NCBI • Google Scholar
19. Brook DW, Rubenstone E, Zhang C, Brook JS (2012) Trajectories of Cigarette Smoking in Adulthood Predict Insomnia among Women in Late Mid-Life. *Sleep Medicine* 13: 1130–1137. pmid:22901402
View Article • PubMed/NCBI • Google Scholar
20. Caldeira KM, O'Grady KE, Garnier-Dykstra LM, Vincent KB, Pickworth WB, Arria AM (2012) Cigarette Smoking among College Students: Longitudinal Trajectories and Health Outcomes. *Nicotine & Tobacco Research* 14: 777–785.
View Article • PubMed/NCBI • Google Scholar
21. Lessov-Schlaggar CN, Kristjansson SD, Bucholz KK, Heath AC, Madden PA (2012) Genetic Influences on Developmental Smoking Trajectories. *Addiction* 107: 1696–1704. pmid:22385035
View Article • PubMed/NCBI • Google Scholar
22. Scott-Sheldon LA, Carey MP, Senn TE, Vanable PA (2010) Smoking Behavior among Low-Income Black Adults: Patterns and Correlates of Smoking Trajectories. *Nicotine & Tobacco Research* 12: 1019–1028.
View Article • PubMed/NCBI • Google Scholar
23. Frosch ZA, Dierker LC, Rose JS, Waldinger RJ (2009) Smoking Trajectories, Health, and Mortality across the Adult Lifespan. *Addictive Behaviors* 34: 701–704. pmid:19428188
View Article • PubMed/NCBI • Google Scholar
24. Amstadter AB, Resnick HS, Nugent NR, Acierno R, Rheingold AA, Minhinnett R, et al. (2009) Longitudinal Trajectories of Cigarette Smoking Following Rape. *Journal of Traumatic Stress* 22: 113–121. pmid:19370699
View Article • PubMed/NCBI • Google Scholar
25. Munafo MR, Heron J, Araya R (2008) Smoking Patterns During Pregnancy and Postnatal Period and Depressive Symptoms. *Nicotine & Tobacco Research* 10: 1609–1620.
View Article • PubMed/NCBI • Google Scholar
26. Bernat DH, Erickson DJ, Widome R, Perry CL, Forster JL (2008) Adolescent Smoking Trajectories: Results from a Population-Based Cohort Study. *Journal of Adolescent Health* 43: 334–340. pmid:18809130
View Article • PubMed/NCBI • Google Scholar
27. Colder CR, Flay BR, Segawa E, Hedeker D (2008) Trajectories of Smoking among Freshmen College Students with Prior Smoking History and Risk for Future Smoking: Data from the University Project Tobacco Etiology Research Network (Uptern) Study. *Addiction* 103: 1534–1543. pmid:18783505
View Article • PubMed/NCBI • Google Scholar
28. Tucker JS, Ellickson PL, Orlando M, Klein DJ (2006) Cigarette Smoking from Adolescence to Young Adulthood: Women's Developmental Trajectories and Associates Outcomes. *Womens Health Issues* 16: 30–37. pmid:16487922
View Article • PubMed/NCBI • Google Scholar
29. Abrams L, Simons-Morton B, Haynie DL, Chen R (2005) Psychosocial Predictors of Smoking Trajectories During Middle and High School. *Addiction* 100: 852–861. pmid:15918815
View Article • PubMed/NCBI • Google Scholar
30. Costello DM, Dierker LC, Jones BL, Rose JS (2008) Trajectories of Smoking from Adolescence to Early Adulthood and Their Psychosocial Risk Factors. *Health Psychology* 27: 811–818. pmid:19025277
View Article • PubMed/NCBI • Google Scholar
31. Fuemmeler BF, Lee C, Ranby KW, Clark T, McClernon FJ, Yang C, et al. (2013) Individual- and Community-Level Correlates of Cigarette-Smoking Trajectories from Age 13 to 32 in a U.S. Population-Based Sample. *Drug and Alcohol Dependence* 132: 301–308. pmid:23499056
View Article • PubMed/NCBI • Google Scholar
32. Pollard MS, Tucker JS, Green HD, Kennedy D, Myong-Hyun G (2010) Friendship Networks and Trajectories of Adolescent Tobacco Use. *Addictive Behaviors* 35: 678–685. pmid:20332061
View Article • PubMed/NCBI • Google Scholar
33. Brown AE, Carpenter MJ, Sutfin EL (2011) Occasional Smoking in College: Who, What, When and Why? *Addictive Behaviors* 36: 1199–1204. pmid:21849231
View Article • PubMed/NCBI • Google Scholar

34. Schane RE, Glantz SA, Ling PM (2009) Nondaily and Social Smoking: An Increasingly Prevalent Pattern. *Archives of Internal Medicine* 169: 1742–1744. pmid:19858429
View Article • PubMed/NCBI • Google Scholar
35. U.S. Bureau of Labor Statistics. National Longitudinal Survey of Youth 1997. <https://www.nlsinfo.org/content/cohorts/NLSY97>. (Accessed 14 Aug 2013)
36. Curran PJ, Obeidat K, Losardo D (2010) Twelve Frequently Asked Questions About Growth Curve Modeling. *Journal of Cognition and Development* 11: 121–136. pmid:21743795
View Article • PubMed/NCBI • Google Scholar
37. U.S. Bureau of Labor Statistics. Sample Weights & Design Effects. <https://www.nlsinfo.org/content/cohorts/nlsy97/using-and-understanding-the-data/sample-weights-design-effects>. (Accessed 4 Dec 2015)
38. Tukey JW (1977) *Exploratory Data Analysis*. New York, NY: Pearson.
39. Tukey JW (1962) The Future of Data Analysis. *The Annals of Mathematical Statistics* 33: 1–67.
View Article • PubMed/NCBI • Google Scholar
40. Muthen B (2003) Statistical and Substantive Checking in Growth Mixture Modeling. *Psychological Methods* 8: 369–377. pmid:14596497
View Article • PubMed/NCBI • Google Scholar
41. Bauer DJ, Curran PJ (2003) Distributional Assumptions of Growth Mixture Models: Implications for Overextraction of Latent Trajectory Classes. *Psychological Methods* 8: 338. pmid:14596495
View Article • PubMed/NCBI • Google Scholar
42. Glenn N (2009) Is the Apparent U-Shape of Well-Being over the Life Course a Result of Inappropriate Use of Control Variables? A Commentary on Blanchflower and Oswald (66: 8, 2008, 1733–1749). *Social Science & Medicine* 69: 481–485; discussion 486–488.
View Article • PubMed/NCBI • Google Scholar
43. Berwick D, Murphy J, Goldman P, Ware J Jr, Barsky A, Weinstein M (1991) Performance of a Five-Item Mental Health Screening Test. *Medical Care* 29: 169–176. pmid:1994148
View Article • PubMed/NCBI • Google Scholar
44. Muthen B, Muthen LK (2000) Integrating Person-Centered and Variable-Centered Analyses: Growth Mixture Modeling with Latent Trajectory Classes. *Alcoholism: Clinical and Experimental Research* 24: 882–891.
View Article • PubMed/NCBI • Google Scholar
45. Nagin DS (1999) Analyzing Development Trajectories: A Semiparametric, Group-Based Approach. *Psychological Methods* 4: 139–157.
View Article • PubMed/NCBI • Google Scholar
46. Muthen LK, Muthen BO (2010) *Mplus User's Guide*. Los Angeles, CA: Muthen & Muthen.
47. Jung T, Wickrama KAS (2008) An Introduction to Latent Class Growth Analysis and Growth Mixture Modeling. *Social and Personality Psychology Compass* 2: 302–317.
View Article • PubMed/NCBI • Google Scholar
48. Long J, Freese J (2006) *Regression Models for Categorical Dependent Variables Using Stata*. 2nd Ed. College Station, TX: Stata Press.
49. Berlin KS, Parra GR, Williams NA (2014) An Introduction to Latent Variable Mixture Modeling (Part 2): Longitudinal Latent Class Growth Analysis and Growth Mixture Models. *J Pediatr Psychol* 39: 188–203. pmid:24277770
View Article • PubMed/NCBI • Google Scholar
50. Muthen B (2016) Re: License Sabc70073114. Personal communication to Lisha N. 17 Nov 2016.
51. Muthen LK. Goodness of Fit Statistics for Mixture Models. <http://www.statmodel.com/discussion/messages/13/119.html?1453602211>. (Accessed 19 Jan 2016)
52. Morgan GB (2014) Mixed Mode Latent Class Analysis: An Examination of Fit Index Performance for Classification. *Structural Equation Modeling: A Multidisciplinary Journal* 22: 76–86.
View Article • PubMed/NCBI • Google Scholar
53. Nagin D (2005) *Group-Based Modeling of Development*. Cambridge, MA: Harvard University Press.
54. Glantz SA (2012) *Primer of Biostatistics*. New York, NY: McGraw Hill.
55. Husten CG (2009) How Should We Define Light or Intermittent Smoking? Does It Matter? *Nicotine & Tobacco Research* 11: 111–121.
View Article • PubMed/NCBI • Google Scholar

56. Cheng KW, Okechukwu CA, McMillen R, Glantz SA (2015) Association between Clean Indoor Air Laws and Voluntary Smokefree Rules in Homes and Cars. *Tobacco Control* 24: 168–174. pmid:24114562
View Article • PubMed/NCBI • Google Scholar
57. Vijayaraghavan M, Messer K, White MM, Pierce JP (2013) The Effectiveness of Cigarette Price and Smoke-Free Homes on Low-Income Smokers in the United States. *American Journal of Public Health* 103: 2276–2283. pmid:24134354
View Article • PubMed/NCBI • Google Scholar
58. Song AV, Dutra LM, Neilands TB, Glantz SA (2015) Association of Smoke-Free Laws with Lower Percentages of New and Current Smokers among Adolescents and Young Adults. *JAMA Pediatrics* 169: e152285. pmid:26348866
View Article • PubMed/NCBI • Google Scholar
59. Cheng KW, Glantz SA, Lightwood JM (2011) Association between Smokefree Laws and Voluntary Smokefree-Home Rules. *American Journal of Preventive Medicine* 41: 566–572. pmid:22099232
View Article • PubMed/NCBI • Google Scholar
60. Hyland A, Higbee C, Travers MJ, Van Deusen A, Bansal-Travers M, King B, et al. (2009) Smoke-Free Homes and Smoking Cessation and Relapse in a Longitudinal Population of Adults. *Nicotine & Tobacco Research* 11: 614–618.
View Article • PubMed/NCBI • Google Scholar
61. American Academy of Pediatrics (2015) Public Policy to Protect Children from Tobacco, Nicotine, and Tobacco Smoke. *Pediatrics* 136: 998–1007. pmid:26504133
View Article • PubMed/NCBI • Google Scholar
62. Song AV, Ling PM, Neilands TB, Glantz SA (2007) Smoking in Movies and Increased Smoking among Young Adults. *American Journal of Preventive Medicine* 33: 396–403. pmid:17950405
View Article • PubMed/NCBI • Google Scholar
63. Shmueli D, Prochaska JJ, Glantz SA (2010) Effect of Smoking Scenes in Films on Immediate Smoking: A Randomized Controlled Study. *American Journal of Preventive Medicine* 38: 351–358. pmid:20307802
View Article • PubMed/NCBI • Google Scholar
64. Ling PM, Lee YO, Hong J, Neilands TB, Jordan JW, Glantz SA (2014) Social Branding to Decrease Smoking among Young Adults in Bars. *American Journal of Public Health* 104: 751–760. pmid:24524502
View Article • PubMed/NCBI • Google Scholar
65. Our Mission. Truth Initiative <http://truthinitiative.org/our-mission>. (Accessed 22 Feb 2016)
66. Centers for Disease Control and Prevention. Campaign Overview. U.S. Department of Health & Human Services December 15, 2015 <http://www.cdc.gov/tobacco/campaign/tips/about/campaign-overview.html>. (Accessed 22 Feb 2016)
67. U.S. Food and Drug Administration. The Real Cost Campaign. FDA 10/21/2015 <http://www.fda.gov/TobaccoProducts/PublicHealthEducation/PublicEducationCampaigns/TheRealCostCampaign/default.htm>. (Accessed 15 Dec 2015)
68. Landman A, Ling PM, Glantz SA (2002) Tobacco Industry Youth Smoking Prevention Programs: Protecting the Industry and Hurting Tobacco Control. *American Journal of Public Health* 92: 917–930. pmid:12036777
View Article • PubMed/NCBI • Google Scholar
69. Ling PM, Glantz SA (2002) Using Tobacco-Industry Marketing Research to Design More Effective Tobacco-Control Campaigns. *JAMA* 287: 2983–2989. pmid:12052128
View Article • PubMed/NCBI • Google Scholar